

## Positive Flavocide® Efficacy Results Presented at Leading Conference on Tick-borne Diseases

### Highlights

- **Purdue University presents new Flavocide® efficacy results for tick control at the 17<sup>th</sup> International Conference on Lyme Borreliosis and Other Tick-Borne Diseases in the USA**
- **Laboratory efficacy studies confirm >90% mortality of *Ixodes scapularis* nymphs, a major vector of Lyme disease in the United States**
- **Residual activity in a prototype formulation of Flavocide was observed for up to four weeks post-application under laboratory conditions**
- **Findings confirm Flavocide's potential for use as a novel acaricide<sup>1</sup> against ticks and mites in public health and animal health applications**

**Bio-Gene Technology Limited (ASX: BGT, Bio-Gene or the Company)**, an Australian company developing the next generation of novel insecticides derived from nature, is pleased to announce that researchers from Purdue University, Indiana, USA, have presented promising new data on Flavocide at the 17<sup>th</sup> International Conference on Lyme Borreliosis and Other Tick-Borne Diseases in Chicago, USA.

This conference is the major international conference on Lyme disease and other tick-borne infections and brings together the most knowledgeable investigators in the world to share scientific discoveries relevant to infections transmitted by *Ixodes* ticks<sup>2</sup>. The conference is arranged by the Mayo Clinic, Michigan State University, and University of North Dakota.

The presentation by researchers Maria Murgia, Phurchhoki Sherpa and Professor Catherine Hill from the Department of Entomology at Purdue University titled 'Assessment of New, Natural Product Formulation to Control Ticks in the Midwest, United States' describes laboratory studies showing that Flavocide formulated in a prototype emulsion in water formulation achieved greater than 90% mortality of *Ixodes scapularis* nymphs within 24 hours of exposure, validating its acaricidal potential. Residual activity was observed for up to four weeks under controlled laboratory conditions.

The research was conducted using a novel test arena, 'Phenocosm' tick assay system that is designed to simulate natural tick habitats such as mulch, leaf litter, and turfgrass. These studies confirm that Flavocide has potential utility in addressing tick-borne diseases of major concern in the United States and globally.

---

<sup>1</sup> *Acaricides* are pesticides that kill members of the arachnid subclass *Acari*, which includes ticks and mites.

<sup>2</sup> *Ixodes* is a genus of hard-bodied ticks (family *Ixodidae*). It includes important disease vectors of animals and humans (tick-borne disease), and some species (notably *Ixodes holocyclus*) inject toxins that can cause paralysis. Some ticks in this genus may transmit the pathogenic bacterium *Borrelia burgdorferi* responsible for causing Lyme disease. Additional organisms that may be transmitted by *Ixodes* are parasites from the genus *Babesia*, which cause babesiosis, and bacteria from the related genus *Anaplasma*, which cause anaplasmosis.

These results have been generated under a vector control program funded by the US Center for Disease Control (CDC) co-ordinated by the Midwest Center of Excellence for Vector Borne Disease and research conducted by Purdue University. Under this program the researchers plan to undertake additional field validation of Flavocide's effect on *Ixodes scapularis* nymphs and techniques to observe their behaviour in the presence of Flavocide.

Commenting on the results, **Catherine Hill, Professor of Entomology at Purdue University** said: *"Tick-borne diseases are a growing public health concern, with Lyme disease cases continuing to rise across the United States. Our research demonstrates that Flavocide shows strong acaricidal activity against Ixodes scapularis, the key vector of Lyme disease, with promising residual efficacy. These results highlight its potential as a valuable new tool in integrated tick management strategies."*

**Peter May, Bio-Gene Technology's Executive Director R&D** said: *"We are delighted that Purdue has presented these data at a leading international conference. Flavocide continues to demonstrate versatility across multiple species that are vectors of disease and this strong performance against ticks opens additional pathways in the public health, companion animal, and livestock markets. With the rising incidence of tick-borne disease, there is a clear global need for safer, more effective acaricides."*

These results further support Bio-Gene's strategy to develop Flavocide as a next-generation insecticide/acaricide, leveraging its novel mode of action and favourable safety profile. The presentation slides are attached.

Approved for release on ASX by Bio-Gene Board of Directors.

- ENDS -

**For further information, please contact:**

Bio-Gene Technology Limited:  
E: [bgt.info@bio-gene.com.au](mailto:bgt.info@bio-gene.com.au)

Matthew Wright  
NWR Communications  
E: [matt@nwrcommunications.com.au](mailto:matt@nwrcommunications.com.au)  
M: 0451 896 420

---

### **About Bio-Gene Technology Limited**

Bio-Gene is an Australian company developing novel bio-insecticides to address the global challenges of insecticide resistance. Its unique products are based on a naturally occurring class of compounds proven to overcome resistance to control pests with minimal impact on human health and the environment.

Bio-Gene's products have multiple applications across crop protection, grain storage, public health and consumer uses. They provide new options derived from nature to meet market demand for effective and safe pest management solutions.

[www.bio-gene.com.au](http://www.bio-gene.com.au)

Flavocide® and Qcide® are registered trademarks of Bio-Gene Technology Limited.

# Assessment of New, Natural Product Formulation to Control Ticks in the Midwest, United States

---

Maria V. Murgia, **Phurchhoki Sherpa**, Catherine A. Hill  
Department of Entomology, Purdue University

17<sup>th</sup> Lyme Borreliosis International Conference, Chicago, September 9, 2025

ersonal use only


# Outline

---

- Background
- Development of the Phenocosm Test Arena
- Product Efficacy Studies
- Tick Biology Studies
- Take Home Messages
- Acknowledgements

# Tick-Borne Diseases, an Increasing Public Health Threat

## Vector Expansion (*Ixodes scapularis*)



Eisen and Eisen, Ticks Tick-borne Dis., 1(10), 2023

**PREVENT TICK BITES!**

- WEAR REPELLENT
- CHECK FOR TICKS DAILY
- SHOWER SOON AFTER BEING OUTDOORS
- CALL YOUR DOCTOR IF YOU GET A FEVER OR RASH

For more information: [www.cdc.gov/ticks](http://www.cdc.gov/ticks)

Centers for Disease Control and Prevention  
National Center for Emerging and Zoonotic Infectious Diseases

0278477-A



# Natural Products (NPs): an Alternative for Tick Control

*Journal of Medical Entomology*, 58(2), 2021, 814–820

doi: 10.1093/jme/tjaa248

Advance Access Publication Date: 20 November 2020

Vector Control, Pest Management, Resistance, Repellents

**Synthetic Pyrethroid, Natural Product, and Entomopathogenic Fungal Acaricide Product Formulations for Sustained Early Season Suppression of Host-Seeking *Ixodes scapularis* (Acari: Ixodidae) and *Amblyomma americanum* Nymphs**

Terry L. Schulze<sup>1</sup> and Robert A. Jordan<sup>2,3</sup>

<sup>1</sup>Terry L. Schulze, Ph.D., Inc., 9 Evergreen Court, Perrineville, NJ 08535, <sup>2</sup>Monmouth County Mosquito Control Division, 1901 Wayside Road, Tinton Falls, NJ 07724, and <sup>3</sup>Corresponding author, e-mail: [robert.jordan@co.monmouth.nj.us](mailto:robert.jordan@co.monmouth.nj.us)

*Journal of Medical Entomology*, 58(1), 2021, 390–397

doi: 10.1093/jme/tjaa188

Advance Access Publication Date: 12 October 2020

Vector Control, Pest Management, Resistance, Repellents

Research

**Evaluating the Effects of Minimal Risk Natural Products for Control of the Tick, *Ixodes scapularis* (Acari: Ixodidae)**

Megan C. Dyer<sup>1,2,5</sup> Matthew D. Requistina<sup>1</sup> Kathryn A. Berger<sup>1,3</sup> Gavino Puggioni<sup>4</sup> and Thomas N. Mather<sup>1,2</sup>

scientific reports

OPEN

**Balsam fir (*Abies balsamea*) needles and their essential oil kill overwintering ticks (*Ixodes scapularis*) at cold temperatures**

Shelley A. Adamo<sup>1,2</sup>, Amal El Nabbout<sup>1</sup>, Laura V. Ferguson<sup>1,2</sup>, Jeffrey S. Zbarsky<sup>1</sup> & Nicoletta Faraone<sup>3</sup>

*Acta Parasitologica* (2024) 69:1141–1147  
<https://doi.org/10.1007/s11686-024-00835-w>

ORIGINAL PAPER

**Chemical Composition and Acaricidal Activity of Essential Oil of *Lavandula dentata* L. on Engorged Females of *Rhipicephalus (Boophilus) microplus* (Acari: Ixodidae)**

Carla Xavier<sup>1</sup> · Eduardo Henrique Amorim Silva<sup>1</sup> · Ivaldo Victor Mota de Siqueira<sup>1</sup> · Ricardo<sup>1</sup> · Vanderson Barbosa Bernardo<sup>2</sup> · Henrique Fonseca Goulart<sup>3</sup> · Julart Santana<sup>3</sup> · Rafael Antonio Nascimento Ramos<sup>1</sup> · Pedro Gregório Vieira Aquino<sup>4</sup> · Carvalho<sup>1</sup>



*Experimental and Applied Acarology* (2022) 86:299–312  
<https://doi.org/10.1007/s10493-022-00692-0>

**Repellency and toxicity of a CO<sub>2</sub>-derived cedarwood oil on hard tick species (Ixodidae)**

Lina B. Flor-Weiler<sup>1</sup> · Robert W. Behle<sup>1</sup> · Fred J. Eller<sup>2</sup> · Ephantus J. Muturi<sup>1</sup> · Alejandro P. Rooney<sup>3</sup>

# Project Goal: Evaluate novel NP actives for potential to control disease vector ticks, tick-bite prevention

**Test Articles:** Active Ingredients/NP formulations; positive controls (bifenthrin/botanical oils); negative control (vehicle)

**Tick Species:** *Ixodes scapularis* nymphs

## Project Roadmap



# Developing The Phenocosm

## ➤ Laboratory and Small-Cage Studies

### ➤ Multiple Substrates:

Filter Paper

Simulated Tick Habitats:

Leaf litter, Mulch, Turfgrass

### ➤ Multipurpose:

- Insecticide Efficacy: ✓ Contact Toxicity  
✓ Repellency  
✓ Residual Activity
- Tick Biology and Behavior



# Product Efficacy Assay Design

- **Ticks:** *Ixodes scapularis* nymphs
- 3 Technical Replicates/Biological Replicate/Product; 5 ticks/Technical Replicate
- **Test Chemistry:** Flavocide™ Oil-in-Water Emulsion (**FLC-EW**; 50% Flavesone; used undiluted)
- **Commercial Products: Talstar-P** (7.9% bifenthrin; used at 0.062% active ingredient);
- Outdoor Pest Control** (OPC; Wondercide; 40.42% Cedarwood oil, 8.74% Sesame oil, 3.5% Sodium Lauryl Sulfate and 47.3% of inert ingredients; used at 0.4 fl. oz./gallon)
- **Negative Control:** untreated filter paper




## Flavocide™

- ❖ “Nature-Identical beta-triketone molecule (flavesone) produced synthetically by a chemical process.”
- ❖ Natural plant compound found in leaves of some Myrtaceae plants (Eucalyptus).
- ❖ New Mode of Action (studies by Pacific Discovery Services).

# Method




## Preparation




in fume hood

## Tick Exposure



in fume hood

## Scoring




on lab bench

# FLC-EW Meets the EPA Registration Targets (>90% mortality at 24hrs PE)




### Predicted Knockdown at 1-hr Post-Exposure



N=3 Biological Replicates

### Model Predicted Probability of Mortality




➤ **OPC: No knockdown or toxicity**

# OPC Exhibits Repellency



### Predicted Probability of Ticks in Contact with the Product Treated Filter Paper




Personal use only

# FLC-EW Exhibits Residual Activity up to 4 Weeks Post-Application



## Model Predicted Probability of Mortality



N = 3 Biological Replicates

ersonal use only

# Summary: Product Efficacy Studies

---

- The Phenocosm enabled the assessment of toxicity, repellency, and residual activity.
- Confirmed Talstar-P contact toxicity (selected as positive control for small-cage studies).
- OPC elicited repellency, but not toxicity under the conditions of this assay.
- FLC-EW met **EPA targets** under laboratory conditions.
- FLC-EW maintained efficacy for up to 4 weeks post-application.
- **FLC-EW is a promising candidate for small-cage trials.**

# Phenocosm Modifications for Simulated Environments (Without Products)

## Phenocosm Modifications



## Tick Labeling and Identification




Photo Credits: M. Murgia, P. Sherpa

Photo Credits: T. Campbell

ersonal use only

DO NOT COPY

DO NOT COPY

# Tick Viability Assay Design

---


- **Ticks: *Ixodes scapularis* nymphs**
- 3 Technical Replicates/Biological Replicate/Substrate; 5 ticks/Technical Replicate
- Location of ticks in the phenocosms
- Viability assessment for up to 4 weeks

ersonal use only

DO NOT COPY


# Phenocosm Supports Tick Viability in Simulated Environments

## Weekly Tick Viability



# Most Ticks Were Found on Substrates

Location of ticks in the phenocosms with substrate




Personal use only

# Take Home Messages

---

- The Phenocosm is a versatile, multipurpose test arena.
- FLC-EW is a promising candidate for small-cage trials.
- Acaricide application may require substrate-disturbing methods.



# Acknowledgements

---

## **MCEVBD**

Dr. Lyric C. Bartholomay  
Dr. Susan Paskewitz

## **CDC**

Dr. Lars Eisen

## **Purdue University**

Dr. Andres F. Oliveros-Diaz  
Dr. Krystal R. Hans  
Dr. Douglas Richmond

## **ENTM Capstone Students**

Brianna L. Gast  
Kayla Bliven

**Purdue University  
Center for Urban and  
Industrial Pest  
Management**

**Bio-Gene Technology**  
Peter May  
Dr. James Wade

**Funding:** MCEVBD, Purdue Internal Funding, BGT